

## **Validation of Satellite Data with Climate Monitoring Potential by Radio Occultation Reference Datasets**

**M. Schwaerz**<sup>1</sup>, G. Kirchengast<sup>1</sup>, A. Leuprecht<sup>1</sup>, J. M. Fritzer<sup>1</sup>, B. Scherllin-Pirscher<sup>1</sup>, and C. Retscher<sup>2</sup>

<sup>1</sup>*Wegener Center for Climate and Global Change (WEGC) and Institute for Geophysics, Astrophysics, and Meteorology/Inst. of Physics (IGAM/IP), University of Graz, Graz, Austria*

<sup>2</sup>*EUMETSAT, Darmstadt, Germany*

marc.schwaerz@uni-graz.at

In the ESA/ESRIN atmospheric Cal/Val context a project of the WEGC prepares and provides long-term radio occultation (RO) reference data from a variety of RO missions (in total up to about 2500 profiles per day) in Generic Earth Observation Metadata Standard (GEOMS) format. Applications of the data include use in the long-loop monitoring of trends related to other spaceborne instruments but also to climate change and variability, in routine validation of atmospheric satellite products, in the validation of geophysical retrieval algorithms, in bridging temporally separated space missions, and in scientific evaluation of atmospheric processes. The project ensures the provision of correlative RO data suitable for in-depth examination of tropospheric and stratospheric fundamental state profiles, such as of temperature and pressure as function of altitude, from satellite observations. This undertaking is highly worthwhile since the unique combination of global coverage, high accuracy and vertical resolution, long-term stability, and virtual all-weather capability makes, in the free atmosphere, the validation with RO data preferable to other methods.

The presentation will introduce the project on multi-mission validation by RO, its setup at the WEGC, and the quality and convenient availability of the RO datasets. It will then focus on the discussion of example results of the multi-year validation of temperature, pressure, and density profiles over the upper troposphere and lower stratosphere (UTLS) from Envisat MIPAS and GOMOS data against collocated RO data from CHAMP, Formosat-3/COSMIC, and MetOp-A/GRAS. This provides valuable hints for future improvements in the processing of the Envisat data. In addition, the RO dataset is inter-validated with profiling data from ground-based sites defined in the Multi-TASTE and VALID projects (Lidars, etc.), which are related to Envisat validation, and against a high-quality radiosonde dataset (RAOBCORE) available from the Univ. of Vienna, Austria. We show how these results help to obtain quantitative estimates on the quality of RO data (e.g., systematic error bounds) and on their utility to serve as reference data for the targeted climate monitoring and validation applications.